

Chapter 1. Rational and Irrational Numbers

Exercise 1.1

Solution (1)

Given rational numbers are $\frac{2}{9}$ and $\frac{3}{8}$

LCM of denominators 9 and 8 is 72

Equivalent fractions of ' $\frac{2}{9}$ ' and ' $\frac{3}{8}$ ' with denominator 72

$$\frac{2}{9} = \frac{2 \times 8}{9 \times 8} = \frac{16}{72}$$

$$\frac{3}{8} = \frac{3 \times 9}{8 \times 9} = \frac{27}{72}$$

Since, $16 < 27$, $\frac{16}{72} < \frac{27}{72}$

So, $\frac{2}{9} < \frac{3}{8}$

A rational number between $\frac{2}{9}$ and $\frac{3}{8}$ is

$$\begin{aligned} & \frac{\frac{2}{9} + \frac{3}{8}}{2} \\ &= \frac{\frac{2 \times 8 + 3 \times 9}{72}}{2} \end{aligned}$$

$$= \frac{16 + 27}{72 \times 2}$$

$$= \frac{43}{144}$$

Descending order of the numbers is $\frac{3}{8}, \frac{43}{144}, \frac{2}{9}$

$$\frac{2}{9} < \frac{43}{144} < \frac{3}{8}$$

Solution (2):

Method II:

L.C.M of 3 and 4 is 12.

Rational number between $\frac{1}{3}$ and $\frac{1}{4}$ is $\frac{\frac{1}{3} + \frac{1}{4}}{2}$

$$= \frac{\frac{4+3}{12}}{2}$$

$$= \frac{7}{12 \times 2}$$

$$= \frac{7}{24}$$

$$\frac{1}{3} = \frac{1 \times 4}{3 \times 4} = \frac{4}{12}$$

$$\frac{1}{4} = \frac{1 \times 3}{4 \times 3} = \frac{3}{12}$$

Since, $4 > 3$, $\frac{4}{12} > \frac{3}{12}$

$\frac{1}{3} > \frac{7}{24}$

$\frac{7}{24}$ is in between $\frac{1}{3}$ and $\frac{1}{4}$. So, $\frac{1}{3} > \frac{7}{24} > \frac{1}{4} \rightarrow \textcircled{1}$

Rational number between $\frac{1}{3}$ and $\frac{7}{24}$ is $\frac{\frac{1}{3} + \frac{7}{24}}{2}$

$$= \frac{\frac{1 \times 8 + 7 \times 1}{24}}{2}$$

$$= \frac{15}{2 \times 24}$$

$$= \frac{15}{48}$$

$\frac{15}{48}$ is in between $\frac{1}{3}$ and $\frac{7}{24}$. So, $\frac{1}{3} > \frac{15}{48} > \frac{7}{24} \rightarrow \textcircled{2}$

From $\textcircled{1}$ and $\textcircled{2}$, ascending order of numbers (increasing order)

$$\text{is } \frac{1}{4} < \frac{7}{24} < \frac{15}{48} < \frac{1}{3}$$

Solution (3)

Given rational numbers are $-\frac{1}{3}$ and $-\frac{1}{2}$

LCM of 3 and 2 is 6.

$$-\frac{1}{3} = \frac{-1 \times 2}{3 \times 2} = \frac{-2}{6} ; \quad -\frac{1}{2} = \frac{-1 \times 3}{2 \times 3} = \frac{-3}{6}$$

Since, $2 < 3$

$$-2 > -3$$

$$\frac{-2}{6} > \frac{-3}{6}$$

$$\text{So, } -\frac{1}{3} > -\frac{1}{2}$$

Rational number between $-\frac{1}{3}$ and $-\frac{1}{2}$ is $\frac{-\frac{1}{3} + (-\frac{1}{2})}{2}$

$$= \frac{-1 \times 2 + (-1) \times 3}{6}$$

$$= \frac{-2 - 3}{6 \times 2}$$

$$= \frac{-5}{12}$$

$$\therefore -\frac{1}{3} > -\frac{5}{12} > -\frac{1}{2} \rightarrow \textcircled{1}$$

Rational number between $-\frac{1}{3}$ and $-\frac{5}{12}$ is $\frac{-\frac{1}{3} + (-\frac{5}{12})}{2}$

$$= \frac{-1 \times 4 + (-5) \times 1}{12}$$

$$= \frac{-4 - 5}{12 \times 2}$$

$$= \frac{-9}{24}$$

$$\therefore -\frac{1}{3} > -\frac{9}{24} > -\frac{5}{12} \rightarrow \textcircled{2}$$

From (1) and (2), $-\frac{1}{3} > -\frac{9}{24} > -\frac{5}{12} > -\frac{1}{2}$

Ascending order (increasing order) of rational numbers

$$-\frac{1}{2}, -\frac{5}{12}, -\frac{9}{24}, -\frac{1}{3}$$

Solution (4):

LCM of 3 and 5 is 15.

$$\frac{1}{3} = \frac{1 \times 5}{3 \times 5} = \frac{5}{15} \quad ; \quad \frac{4}{5} = \frac{4 \times 3}{5 \times 3} = \frac{12}{15}$$

Since, $5 < 12$

$$\text{So, } \frac{5}{15} < \frac{12}{15}$$

$$\frac{1}{3} < \frac{4}{5}$$



Rational number between $\frac{1}{3}$ and $\frac{4}{5}$ is $\frac{\frac{1}{3} + \frac{4}{5}}{2}$


$$= \frac{\frac{1 \times 5 + 4 \times 3}{3 \times 5}}{2}$$

$$= \frac{\frac{5 + 12}{15}}{2}$$

stud flare

—IGNITE YOUR LEARNING SPARK—


Rational number between $\frac{1}{3}$ and $\frac{17}{30}$ is



$$\frac{\frac{1}{3} + \frac{17}{30}}{2}$$


$$= \frac{\frac{1 \times 10 + 17}{30}}{2}$$

$$= \frac{27}{60}$$



Rational number between $\frac{17}{30}$ and $\frac{4}{5}$ is

$$\begin{aligned} & \frac{\frac{17}{30} + \frac{4}{5}}{2} \\ &= \frac{\frac{17 + 4 \times 6}{30}}{2} \\ &= \frac{41}{60} \end{aligned}$$



→ Rational numbers between $\frac{1}{3}$ and $\frac{4}{5}$ are



—IGNITE YOUR LEARNING SPARK—

Descending order (decreasing order) of numbers are

$$\frac{4}{5}, \frac{41}{60}, \frac{17}{30}, \frac{27}{60}, \frac{1}{3}$$

Solution ⑤:

A rational number between 4 and 4.5 is $\frac{4+4.5}{2} = \frac{8.5}{2}$
 $= 4.25$

A rational number between 4 and 4.25 is $\frac{4+4.25}{2} = \frac{8.25}{2}$
 $= 4.125$

A rational number between 4 and 4.125 is $\frac{4+4.125}{2} = \frac{8.125}{2}$
 $= 4.0625$

Three rational numbers between 4 and 4.5 are

$$4.0625, 4.125, 4.25$$

Solution ⑥:

We need to insert six rational numbers between 3 and 4. So, we multiply both numerator and denominator of rational numbers with $6+1$ i.e. 7.

$$\text{So, } \frac{3}{1} = \frac{3 \times 7}{1 \times 7} = \frac{21}{7}$$

$$\frac{4}{1} = \frac{4 \times 7}{1 \times 7} = \frac{28}{7}$$

We have, $21 < 22 < 23 < 24 < 25 < 26 < 27$

$$\Rightarrow \frac{21}{7} < \frac{22}{7} < \frac{23}{7} < \frac{24}{7} < \frac{25}{7} < \frac{26}{7} < \frac{27}{7} < \frac{28}{7}$$

Therefore, six rational numbers between 3 and 4

are $\frac{22}{7}, \frac{23}{7}, \frac{24}{7}, \frac{25}{7}, \frac{26}{7}, \frac{27}{7}$.

Solution ⑦:

We need to insert five rational numbers between $\frac{3}{5}$ and $\frac{4}{5}$. So, we multiply both numerator and denominator with $5+1$ i.e. 6.

$$\text{So, } \frac{3}{5} = \frac{3 \times 6}{5 \times 6} = \frac{18}{30}$$

$$\frac{4}{5} = \frac{4 \times 6}{5 \times 6} = \frac{24}{30}$$

We have, $18 < 19 < 20 < 21 < 22 < 23 < 24$

$$\Rightarrow \frac{18}{30} < \frac{19}{30} < \frac{20}{30} < \frac{21}{30} < \frac{22}{30} < \frac{23}{30} < \frac{24}{30}$$

Therefore, five rational numbers between $\frac{3}{5}$ and $\frac{4}{5}$.

$$\frac{19}{30}, \frac{20}{30}, \frac{21}{30}, \frac{22}{30}, \frac{23}{30}$$

Solution (8):

LCM of 5 and 7 is 35.

$$\frac{-2}{5} = \frac{-2 \times 7}{5 \times 7} = \frac{-14}{35} \quad ; \quad \frac{1}{7} = \frac{1 \times 5}{7 \times 5} = \frac{5}{35}$$

We need to insert ten rational numbers between $\frac{-2}{5}$ ($= \frac{-14}{35}$) and $\frac{1}{7}$ ($= \frac{5}{35}$). So, we can select any ten numbers between -14 and 5 as numerators and '35' as denominator.

$$\therefore \frac{-13}{35}, \frac{-12}{35}, \frac{-11}{35}, \frac{-10}{35}, \frac{-9}{35}, \frac{-8}{35}, \frac{-7}{35}, \frac{-6}{35}$$

$$\frac{-5}{35}, \frac{-4}{35} \text{ are ten rational numbers}$$

which are in between $\frac{-2}{5}$ and $\frac{1}{7}$.

Solution (9)

LCM of 2 and 3 is 6.

$$\frac{1}{2} = \frac{1 \times 3}{2 \times 3} = \frac{3}{6} \quad ; \quad \frac{2}{3} = \frac{2 \times 2}{3 \times 2} = \frac{4}{6}$$

We need to insert six rational numbers. So, multiply both numerator and denominator by 6+1 i.e., 7.

$$\frac{3}{6} = \frac{3 \times 7}{6 \times 7} = \frac{21}{42} \quad ;$$

$$\frac{4}{6} = \frac{4 \times 7}{6 \times 7} = \frac{28}{42}$$

Since, $21 < 22 < 23 < 24 < 25 < 26 < 27 < 28$

$$\frac{21}{42} < \frac{22}{42} < \frac{23}{42} < \frac{24}{42} < \frac{25}{42} < \frac{26}{42} < \frac{27}{42} < \frac{28}{42}$$

Therefore, Six numbers between $\frac{1}{2}$ and $\frac{2}{3}$

$$\text{are } \frac{22}{42}, \frac{23}{42}, \frac{24}{42}, \frac{25}{42}, \frac{26}{42}, \frac{27}{42}$$

Exercise - 1.2

Solution 1:

Let $\sqrt{5}$ be a rational number, then

$\sqrt{5} = \frac{p}{q}$, where p, q are integers, $q \neq 0$ and p, q have no common factors (except 1).

$$\Rightarrow 5 = \frac{p^2}{q^2}$$

$$p^2 = 5 \cdot q^2 \quad \longrightarrow (i)$$

As '5' divides $5q^2$, so '5' divides p^2 and '5' is prime

$\Rightarrow 5$ divides p .

Let $p = 5 \cdot m$, where m is an integer

Substituting the value of 'p' in (i)

$$(5m)^2 = 5 \cdot q^2$$

$$25 \cdot m^2 = 5 \cdot q^2$$

$$\Rightarrow q^2 = 5 \cdot m^2$$

As 5 divides $5m^2$, so 5 divides q^2 but 5 is prime

$\Rightarrow 5$ divides q .

Thus, p and q have a common factor 5. This contradicts that 'p' and 'q' have no common factors (except 1).

Hence, $\sqrt{5}$ is not rational number.

So, we conclude $\sqrt{5}$ is an irrational number.

Solution 2:

Let $\sqrt{7}$ be a rational number.

$\sqrt{7} = \frac{p}{q}$, where p, q are integers, $q \neq 0$ and p, q have no common factors (except 1).

$$\Rightarrow 7 = \frac{p^2}{q^2}$$

$$p^2 = 7 \cdot q^2 \rightarrow (i)$$

As 7 divides $7q^2$, so 7 divides p^2 and 7 is a prime

\Rightarrow 7 divides p

Let $p = 7m$, where 'm' is an integer

Substitute this value of 'p' in \therefore we have.

$$(7m)^2 = 7 \cdot q^2$$

$$49 \cdot m^2 = 7 \cdot q^2$$

$$7m^2 = q^2$$

As 7 divides $7m^2$, so 7 divides q^2 but 7 is a prime number.

\Rightarrow 7 divides q

Thus, 'p' and q have a common factor 7. This contradicts that p and q have no common factor (except 1).

Hence, $\sqrt{7}$ is not a rational number.

So, we conclude $\sqrt{7}$ is irrational number.

Solution 3:

Let $\sqrt{6}$ be a rational number.

$\sqrt{6} = \frac{p}{q}$, where p and q are integers, $q \neq 0$ and p, q have no common factors (except 1).

$$\Rightarrow 6 = \frac{p^2}{q^2}$$

$$p^2 = 6 \cdot q^2 \rightarrow (i)$$

As '2' divides $6q^2$, so 2 divides p^2 but 2 is prime

\Rightarrow 2 divides p .

Let $p = 2 \cdot m$, where 'm' is an integer.

Substitute this value of 'p' in (i)

$$(2m)^2 = 6 \cdot q^2$$

$$4 \cdot m^2 = 6 \cdot q^2$$

$$2 \cdot m^2 = 3 \cdot q^2$$

'2' divides '2m²', so 2 divides '3q²'

2 should either divide '3' or divide q².

But 2 should does not divide 3.

Therefore, 2 divides q² and 2 is a prime

2 divides q.

Thus, p and q have a common factor 2. This contradicts that 'p' and 'q' have no common factors (except 1).

Hence, $\sqrt{6}$ is not rational number.

So, we conclude $\sqrt{6}$ is irrational number.

Solution 4 :-

Let $\frac{1}{\sqrt{11}}$ be a rational number.

$\frac{1}{\sqrt{11}} = \frac{p}{q}$, where 'p', 'q' are integers, $q \neq 0$ and p, q have no common factors (except 1).

$$\Rightarrow \frac{1}{11} = \frac{p^2}{q^2}$$

$$\Rightarrow q^2 = 11 \cdot p^2 \rightarrow \text{ii}$$

As 11 divides $11p^2$, so 11 divides q^2 but 11 is a prime.

$$\Rightarrow 11 \text{ divides } q.$$

$q = 11m$, where 'm' is an integer.

$$(11m)^2 = 11p^2$$

$$\Rightarrow 121 \cdot m^2 = 11 \cdot p^2$$

$$\Rightarrow p^2 = 11 \cdot m^2$$

As 11 divides $11m^2$, so 11 divides p^2 but 11 is a prime.

$$\Rightarrow 11 \text{ divides } p.$$

Thus, p and q have a common factor 11. This contradicts the fact that p and q has no common factors (except 1).

Hence, $\frac{1}{\sqrt{11}}$ is not rational number.

So, we conclude $\frac{1}{\sqrt{11}}$ is irrational number.

Solution 5:

Let $\sqrt{2}$ is a rational number.

$\sqrt{2} = \frac{p}{q}$, where p and q are integers, $q \neq 0$ and p, q have no common factors (except 1).

$$\Rightarrow 2 = \frac{p^2}{q^2}$$

$$p^2 = 2 \cdot q^2 \rightarrow (i)$$

As '2' divides $2q^2$, so 2 divides p^2 but 2 is prime.

\Rightarrow 2 divides p .

Let $p = 2m$, where 'm' is an integer.

Substitute this value of p in (i).

$$(2m)^2 = 2q^2$$

$$4m^2 = 2q^2$$

$$\Rightarrow q^2 = 2m^2$$

As 2 divides $2m^2$, so 2 divides q^2 but 2 is prime.

2 divides q

Thus, p and q have a common factor 2. This contradicts the fact that p and q has no common factor (except 1).

Hence, $\sqrt{2}$ is not rational number.

So, we conclude $\sqrt{2}$ is irrational number.

Let us assume $3 - \sqrt{2}$ is rational number, say r .

$$\text{Thus, } 3 - \sqrt{2} = r \Rightarrow 3 - r = \sqrt{2}$$

As 'r' is rational, $3 - r$ is rational $\Rightarrow \sqrt{2}$ is rational

this contradicts the fact that $\sqrt{2}$ is irrational

Hence, our assumption is wrong. Therefore, $3 - \sqrt{2}$ is an irrational number.

Solution 6:

Let $\sqrt{3}$ is a rational number.

$\sqrt{3} = \frac{p}{q}$, where p and q are integers, $q \neq 0$ and p, q have no common factors (except 1).

$$3 = \frac{p^2}{q^2}$$

$$\Rightarrow p^2 = 3q^2 \longrightarrow (i)$$

As '3' divides $3q^2$, so 3 divides p^2 but 3 is a prime.

$$\Rightarrow 3 \text{ divides } p$$

Let $p = 3 \cdot m$, where 'm' is an integer.

substituting this value of p in (i).

$$(3m)^2 = 3q^2$$

$$9m^2 = 3q^2$$

$$\Rightarrow q^2 = 3m^2$$

As '3' divides $3m^2$, so '3' divides q^2 but 3 is not prime.

$$\Rightarrow 3 \text{ divides } q$$

Thus, p and q have a common factor 3. This contradicts the fact that p and q has no common factor (except 1).

Hence, $\sqrt{3}$ is not rational number.

So, we conclude $\sqrt{3}$ is irrational number.

Let us assume $\frac{2}{5}\sqrt{3}$ is a rational number, say r .

$$\text{Thus } \frac{2}{5}\sqrt{3} = r \Rightarrow \sqrt{3} = \frac{5}{2} \cdot r$$

As r is rational, $\frac{5}{2}r$ is rational $\Rightarrow \sqrt{3}$ is rational.

This contradicts the fact that $\sqrt{3}$ is irrational.

Hence, our assumption is wrong. Therefore, $\frac{2}{5}\sqrt{3}$ is irrational number.

Solution 7:

Let $\sqrt{5}$ is a rational number.

$\sqrt{5} = \frac{p}{q}$, where p and q are integers, $q \neq 0$ and p, q have no common factors (except 1).

$$5 = \frac{p^2}{q^2}$$

$$\Rightarrow p^2 = 5q^2 \rightarrow (i)$$

As 5 divides $5q^2$, so 5 divides p^2 but 5 is a prime.

$$\Rightarrow 5 \text{ divides } p.$$

Let $p = 5m$ where m is an integer.

Substitute this value of m in (i)

$$(5m)^2 = 5q^2$$

$$25m^2 = 5q^2$$

$$q^2 = 5m^2$$

As 5 divides $5m^2$, 5 divides q^2 but 5 is a prime.

5 divides q

Thus, p and q have a common factor 5. This contradicts the fact that p and q has no common factors (except 1).

Hence, $\sqrt{5}$ is not rational number.

So, we conclude $\sqrt{5}$ is irrational number.

Let us assume $-3+2\sqrt{5}$ is a rational number, say r .

$$\text{Thus, } -3+2\sqrt{5} = r \Rightarrow -3-r = 2\sqrt{5}$$

$$\Rightarrow \sqrt{5} = \frac{-(3+r)}{2}$$

As r is rational, $-\left(\frac{3+r}{2}\right)$ is rational $\Rightarrow \sqrt{5}$ is rational.

This contradict the fact that $\sqrt{5}$ is irrational.

Hence, our assumption is wrong, therefore, $-3+2\sqrt{5}$ is irrational number.

Solution (8):

(i) Let $5 + \sqrt{2}$ is rational number, say r .

$$5 + \sqrt{2} = r \Rightarrow \sqrt{2} = r - 5$$

As r is rational, $r - 5$ is rational $\Rightarrow \sqrt{2}$ is rational.

This contradicts the fact that $\sqrt{2}$ is irrational.

Hence, our assumption is wrong. Therefore, $5 + \sqrt{2}$ is an irrational number.

(ii)

Let $3 - 5\sqrt{3}$ is rational number, say r .

$$3 - 5\sqrt{3} = r \Rightarrow 5\sqrt{3} = 3 - r$$

$$\Rightarrow \sqrt{3} = \left(\frac{3-r}{5}\right)$$

As r is rational, $\left(\frac{3-r}{5}\right)$ is rational $\Rightarrow \sqrt{3}$ is rational.

This contradicts the fact that $\sqrt{3}$ is irrational.

Hence, our assumption is wrong. Therefore, $3 - 5\sqrt{3}$ is an irrational number.

(iii)

Let $2\sqrt{3} - 7$ is a rational number, say r .

$$2\sqrt{3} - 7 = r \Rightarrow 2\sqrt{3} = r + 7$$

$$\sqrt{3} = \frac{r+7}{2}$$

As r is rational, $\left(\frac{r+7}{2}\right)$ is rational $\Rightarrow \sqrt{3}$ is rational.

This contradicts the fact that $\sqrt{3}$ is irrational.

Hence, our assumption is wrong. Therefore, $2\sqrt{3} - 7$ is an irrational number.

Solution 8:

(iv) Let $\sqrt{2} + \sqrt{5}$ is a rational number, say r .

$$\sqrt{2} + \sqrt{5} = r$$

$$\sqrt{5} = r - \sqrt{2}$$

$$(\sqrt{5})^2 = (r - \sqrt{2})^2 \quad (\text{on squaring both sides})$$

$$5 = r^2 + (\sqrt{2})^2 - 2 \times r \times \sqrt{2} \quad [\because (a-b)^2 = a^2 + b^2 - 2ab]$$

$$5 = r^2 + 2 - 2\sqrt{2} \cdot r$$

$$2\sqrt{2} \cdot r = r^2 - 3$$

$$\sqrt{2} = \frac{r^2 - 3}{2 \cdot r}$$

As r is rational, $r^2 - 3$ is rational, $\left(\frac{r^2 - 3}{2r}\right)$ is rational
 $\Rightarrow \sqrt{2}$ is rational

this contradicts the fact that $\sqrt{2}$ is irrational.

Hence, our assumption is wrong. Therefore,

$(\sqrt{2} + \sqrt{5})$ is an irrational number.

Exercise 1.3

solution 1:



PQ is a number line.

We have two right angle triangles. They are

$\triangle OAB$ and $\triangle OCD$.

In a right angled triangle,

$$(\text{hypotenuse})^2 = (\text{side 1})^2 + (\text{side 2})^2$$

$$\therefore OB^2 = OA^2 + AB^2$$

$$OB^2 = 3^2 + 1^2 \quad (\because OA = 3, AB = 1)$$

$$OB^2 = 9 + 1$$

$$OB = \sqrt{10}$$

Similarly, in $\triangle OCD$,

$$OD^2 = OC^2 + CD^2$$

$$OD^2 = 4^2 + 1^2$$

$$(\because OC = 4, CD = 1)$$

$$OD^2 = 16 + 1$$

$$OD = \sqrt{17}$$

Solution 2:-

(i) $\frac{36}{100}$

$$\begin{array}{r|l} & 0.036 \\ 100 & 360 \\ & \underline{300} \\ & 600 \\ & \underline{600} \\ & 0 \end{array}$$

Remainder becomes zero.

Decimal expansion of $\frac{36}{100} (=0.36)$ is terminating.

(ii) $4\frac{1}{8}$

$$= \frac{4 \times 8 + 1}{8}$$

$$= \frac{33}{8}$$

$$\begin{array}{r|l} & 4.125 \\ 8 & 33 \\ & \underline{32} \\ & 10 \\ & \underline{8} \\ & 20 \\ & \underline{16} \\ & 40 \\ & \underline{40} \\ & 0 \end{array}$$

Remainder becomes zero

Decimal expansion of $4\frac{1}{8} (=4.125)$ is terminating.

(iii) $\frac{2}{9}$

	0.22
9	20 18
	20 18
	2

← Remainder is repeating

In the above decimal expansion, remainder is repeating. So, it is a non-terminating decimal.

So, $\frac{2}{9} = 0.222\dots = 0.\dot{2} = 0.\bar{2}$

(iv) $\frac{2}{11}$

—IGNITE YOUR LEARNING SPARK—

	0.1818
11	20 11
	90 88
	20 11
	90 88
	2

← Remainder '2' is repeating.

Decimal expansion of $\frac{2}{11} = 0.1818\dots$

Here, remainder is repeating. So, it is a non-terminating repeating decimal.

$\therefore \frac{2}{11} = 0.\overline{18}$

(v) $\frac{3}{13}$

	0.230769
13	30 26
	40 39
	100 91
	90 78
	120 117
	3 ← Remainder '3' is repeating.

Decimal expansion of $\frac{3}{13}$ is 0.230769.....

Here, remainder is repeating. So, it is a non-terminating repeating decimal.

$\therefore \frac{3}{13} = 0.\overline{230769}$

(vi) $\frac{329}{400}$

	0.8225
100	3290 3200
	900 800
	1000 800
	2000 2000
	0 ← Remainder is zero

Decimal Expansion of $\frac{329}{400} = 0.8225$.

∴ It is a ~~repet~~ terminating decimal expansion.

Solution (3) :-

(i) $\frac{13}{3125}$

Prime factorization of denominator 3125.

$$\begin{aligned} 3125 &= 5 \times 5 \times 5 \times 5 \times 5 \times 1 \\ &= 5^5 \times 1 \\ &= 1 \times 5^5 \end{aligned}$$

$$\begin{array}{r} 5 \overline{) 3125} \\ \underline{15625} \\ 5 \overline{) 625} \\ \underline{3125} \\ 5 \overline{) 125} \\ \underline{625} \\ 5 \overline{) 25} \\ \underline{25} \\ 5 \overline{) 5} \\ \underline{5} \\ 1 \end{array}$$

∴ $3125 = 2^0 \times 5^5$ ($\because 2^0 = 1$)

Since, denominator is in the form of $2^0 \times 5^5$,
the decimal expansion of $\frac{13}{3125}$ is terminating.

(ii) $\frac{17}{8}$

Prime factorization of denominator 8.

$$8 = 2 \times 2 \times 2$$

$$8 = 2^3 \times 1$$

$$8 = 2^3 \times 5^0$$

$$\begin{array}{r} 2 \overline{) 8} \\ \underline{4} \\ 2 \overline{) 4} \\ \underline{2} \\ 2 \overline{) 2} \\ \underline{2} \\ 1 \end{array}$$

Since denominator is in the form of $2^3 \times 5^0$,
decimal expansion of $\frac{17}{8}$ is terminating.

$$\underline{\underline{(iii) \frac{23}{75}}}$$

Prime factorization of 75.

$$75 = 3 \times 5 \times 5 \times 1$$

$$75 = 3 \times 5^2 \times 1$$

$$75 = 3 \times 2^0 \times 5^2 \quad (\because 2^0 = 1)$$

$$\begin{array}{r} 3 \overline{) 75} \\ 5 \overline{) 25} \\ 5 \overline{) 5} \\ 1 \end{array}$$

Since, denominator contains prime factor 3 other than 2 or 5.

Decimal expansion of $\frac{23}{75}$ is non-terminating.

$$\underline{\underline{(iv) \frac{6}{15}}}$$

Both numerator and denominator contains common factor 3.

$$\frac{6}{15} = \frac{3 \times 2}{3 \times 5} = \frac{2}{5}$$

$$\therefore \frac{6}{15} = \frac{2}{5}$$

Since, denominator is in the form $2^0 \times 5^1$.

Decimal expansion of $\frac{6}{15} (= \frac{2}{5})$ is terminating.

$$\underline{\underline{(v) \frac{1258}{625}}}$$

Prime factorization of denominator 625.

$$625 = 5 \times 5 \times 5 \times 5 \times 1$$

$$625 = 5^4 \times 2^0$$

$$\begin{array}{r} 5 \overline{) 625} \\ 5 \overline{) 125} \\ 5 \overline{) 25} \\ 5 \overline{) 5} \\ 1 \end{array}$$

Since, denominator is in the form $2^0 \times 5^4$, decimal expansion

of $\frac{1258}{625}$ is terminating.

$$(vi) \frac{77}{210}$$

Both numerator and denominator contains common factor 7.

$$\frac{77}{210} = \frac{7 \times 11}{7 \times 30} = \frac{11}{30}$$

$$\therefore \frac{77}{210} = \frac{11}{30}$$

Prime factorization of denominator 30.

$$30 = 2 \times 3 \times 5 \times 1$$

$$30 = 3 \times 2 \times 5$$

$$\begin{array}{r|l} 2 & 30 \\ \hline 3 & 15 \\ \hline 5 & 5 \\ \hline & 1 \end{array}$$

Since, denominator contains prime factor 3 other 2 or 5

Decimal expansion of $\frac{77}{210}$ is non-terminating.

Solution (4) :-

Expressing both numerator and denominator of fraction

$\frac{987}{10500}$ as product of prime numbers by prime factorization method

$$\begin{array}{r|l} 3 & 987 \\ \hline 7 & 329 \\ \hline 47 & 47 \\ \hline & 1 \end{array}$$

$$\therefore 987 = 3 \times 7 \times 47$$

$$\begin{array}{r|l} 2 & 10500 \\ \hline 2 & 5250 \\ \hline 3 & 2625 \\ \hline 5 & 875 \\ \hline 5 & 175 \\ \hline 5 & 35 \\ \hline & 7 \end{array}$$

$$\therefore 10500 = 2 \times 2 \times 3 \times 5 \times 5 \times 5 \times 7$$

$$\frac{987}{10500} = \frac{\cancel{3} \times \cancel{7} \times 47}{2 \times 2 \times \cancel{3} \times 5 \times 5 \times 5 \times \cancel{7}}$$

$$= \frac{47}{2^2 \times 5^3}$$

Since, denominator is in the form $2^2 \times 5^3$, decimal

Expansion of $\frac{987}{10500}$ is terminating

Solution (5) :-

(i) $\frac{17}{8}$

Prime factorization of denominator 8

$$8 = 2 \times 2 \times 2 \times 1$$

$$8 = 2^3 \times 5^0 \quad (\because a^0 = 1)$$

$$\begin{array}{r|l} 2 & 8 \\ \hline 2 & 4 \\ \hline 2 & 2 \\ \hline & 1 \end{array}$$

$$\frac{17}{8} = \frac{17}{2^3}$$

$$= \frac{17 \times 5^3}{2^3 \times 5^3} \quad (\text{By multiplying both numerator and denominator with } 5^3).$$

$$= \frac{17 \times 125}{(2 \times 5)^3}$$

$$= \frac{2125}{10^3}$$

$$= 2.125$$

(Since, denominator is in the form 10^3 , decimal expansion is obtained by moving decimal point to three digits from right).

$$\therefore \frac{17}{8} = 2.125 //$$

$$\begin{array}{r} 2.125 \\ \times 17 \\ \hline 14375 \\ 42500 \\ \hline 212500 \end{array}$$

$$(ii) \frac{13}{3125}$$

Prime factorization of 3125.

$$3125 = 5 \times 5 \times 5 \times 5 \times 5 = 5^5$$

$$\begin{array}{r} 5 \overline{) 3125} \\ \underline{5} \\ 625 \\ \underline{5} \\ 125 \\ \underline{5} \\ 25 \\ \underline{5} \\ 5 \\ \underline{5} \\ 1 \end{array}$$

$$\frac{13}{3125} = \frac{13}{5^5}$$

$$= \frac{13 \times 2^5}{5^5 \times 2^5} \quad (\text{Multiplying numerator and denominator by } 2^5)$$

$$= \frac{13 \times 32}{(2 \times 5)^5} \quad \left(\begin{array}{l} 2^5 = 2 \times 2 \times 2 \times 2 \times 2 = 32 \\ \& a^m \times b^m = (a \times b)^m \end{array} \right)$$

$$= \frac{416}{10^5}$$

$$= 0.00416$$

$$\begin{array}{r} 32 \\ \times 13 \\ \hline 96 \\ 32 \times \\ \hline 416 \end{array}$$

$$\therefore \frac{13}{3125} = 0.00416$$

$$(iii) \frac{7}{80}$$

Prime factorization of 80.

$$80 = 2 \times 2 \times 2 \times 2 \times 5$$

$$80 = 2^4 \times 5^1$$

$$\begin{array}{r} 2 \overline{) 80} \\ \underline{2} \\ 40 \\ \underline{2} \\ 20 \\ \underline{2} \\ 10 \\ \underline{2} \\ 5 \\ \underline{5} \\ 1 \end{array}$$

$$\frac{7}{80} = \frac{7}{2^4 \times 5^1}$$

$$= \frac{7 \times 5^3}{2^4 \times 5^1 \times 5^3}$$

$$= \frac{7 \times 125}{2^4 \times 5^4}$$

(Multiplying numerator and denominator by 5^3)

$$= \frac{7 \times 125}{(2 \times 5)^4}$$

$$= \frac{875}{10^4}$$

$$= 0.0875$$

$$\begin{array}{r} 125 \\ \times 7 \\ \hline 875 \end{array}$$

(Since, denominator is 10^4 , decimal expansion can be obtained by moving decimal point of numerator to four digits from right.)

$$\therefore \frac{7}{80} = 0.0875$$

(iv) $\frac{6}{15}$

Prime factorization of 6 and 15

$$\begin{array}{r} 2 \overline{) 6} \\ 3 \overline{) 3} \\ \hline \end{array}$$

$$\begin{array}{r} 3 \overline{) 15} \\ 5 \overline{) 5} \\ \hline \end{array}$$

IGNITE YOUR LEARNING SPARK

$$\therefore 6 = 2 \times 3$$

$$\therefore 15 = 3 \times 5$$

$$\frac{6}{15} = \frac{2 \times 3}{3 \times 5} = \frac{2}{5}$$

$$= \frac{2 \times 2}{5 \times 2}$$

$$= \frac{4}{10}$$

$$= 0.4$$

(By multiplying both numerator and denominator by 2.)

$$(v) \frac{2^3 \times 7}{5^4}$$

$$= \frac{2^3 \times 7 \times 2^4}{5^4 \times 2^4}$$

(By multiplying both numerator and denominator by 2^4)

$$= \frac{4 \times 7 \times 16}{(2 \times 5)^4}$$

$$= \frac{28 \times 16}{10^4}$$

$$= \frac{448}{10^4}$$

$$= 0.0448$$

$$\therefore \frac{2^3 \times 7}{5^4} = 0.0448$$

$$\begin{array}{r} 28 \\ \times 16 \\ \hline 168 \\ 28 \times \\ \hline 448 \end{array}$$

$$(vi) \frac{237}{1500}$$

IGNITE YOUR LEARNING SPARK

Prime factorization of 237 and 1500.

$$\begin{array}{r} 3 \overline{)237} \\ \underline{79} \\ 79 \\ \underline{1} \end{array}$$

$$\begin{array}{r} 2 \overline{)1500} \\ \underline{750} \\ 3 \overline{)375} \\ \underline{125} \\ 5 \overline{)25} \\ \underline{5} \\ 5 \overline{)5} \\ \underline{1} \end{array}$$

$$\therefore 237 = 3 \times 79$$

$$\therefore 1500 = 2 \times 2 \times 3 \times 5 \times 5 \times 5$$

$$\frac{237}{1500} = \frac{\cancel{3} \times 79}{2^2 \times \cancel{3} \times 5^3}$$

$$= \frac{79}{2^2 \times 5^3}$$

$$= \frac{79 \times 2}{2^2 \times 5^3 \times 2} \quad \left(\text{Multiplying both numerator and denominator by '2'} \right)$$

$$= \frac{158}{2^3 \times 5^3}$$

$$= \frac{158}{(10)^3}$$

$$= 0.158$$

$$\therefore \frac{237}{1500} = 0.158$$

Solution (6):

Given rational number $\frac{257}{5000}$

Prime factorization of denominator 5000

$$\begin{aligned} \therefore 5000 &= 2 \times 2 \times 2 \times 5 \times 5 \times 5 \times 5 \\ &= 2^3 \times 5^4 \end{aligned}$$

Thus, denominator of rational number is in the form $2^m \times 5^n$, where $m=3$ and $n=4$.

	5000
2	2500
2	1250
5	625
5	125
5	25
5	5
	1

$$\therefore \frac{257}{5000} = \frac{257}{2^3 \times 5^4}$$

$$= \frac{257 \times 2}{2^3 \times 5^4 \times 2}$$

$$= \frac{514}{2^4 \times 5^4}$$

(By multiplying numerator and denominator by 2)

$$= \frac{514}{(2 \times 5)^4}$$

$$= \frac{514}{10^4}$$

$$= 0.0514$$

∴ Decimal expansion of $\frac{257}{5000}$ is 0.0514 .

Solution (7) :-

Decimal expansion of $\frac{1}{7}$:

	0.142857
7	10
	7
	30
	28
	20
	14
	60
	56
	40
	35
	50
	49
	1

← Remainder '1' is repeated.

∴ Decimal Expansion of $\frac{1}{7}$ is non-terminating repeating

$$\frac{1}{7} = 0.\overline{142857}$$

$\frac{2}{7}$ can be written as $2 \times \frac{1}{7}$

$$\begin{aligned}\text{Decimal Expansion of } \frac{2}{7} &= 2 \times \frac{1}{7} \\ &= 2 \times 0.\overline{142857} \\ &= 0.\overline{285714}\end{aligned}$$

Similarly,

$$\begin{aligned}\frac{3}{7} &= 3 \times \frac{1}{7} = 3 \times 0.\overline{142857} \\ &= 0.\overline{428571}\end{aligned}$$

$$\begin{aligned}\frac{4}{7} &= 4 \times \frac{1}{7} = 4 \times 0.\overline{142857} \\ &= 0.\overline{571428}\end{aligned}$$

$$\begin{aligned}\frac{5}{7} &= 5 \times \frac{1}{7} = 5 \times 0.\overline{142857} \\ &= 0.\overline{714285}\end{aligned}$$

$$\begin{aligned}\frac{6}{7} &= 6 \times \frac{1}{7} = 6 \times 0.\overline{142857} \\ &= 0.\overline{857142}\end{aligned}$$

Solution (8) :-

$$(i) \text{ Let } x = 0.\overline{3} = 0.3333\ldots \rightarrow \textcircled{1}$$

As there is one repeating digit after the decimal point. So multiplying both sides of Eq (1) by 10.

$$10x = 3.333\ldots \rightarrow \textcircled{2}$$

Subtracting (1) from (2), we get.

$$10x - x = 3.333\ldots - 0.333\ldots$$

$$9x = 3$$

$$x = \frac{3}{9}$$

$$x = \frac{1}{3}$$

$\therefore x = 0.\bar{3} = \frac{1}{3}$, which is in $\frac{p}{q}$ form.

(ii) Let $x = 5.\bar{2} = 5.222\dots \rightarrow \textcircled{1}$

As there is one repeating digit after the decimal point, so multiplying both sides of Eq $\textcircled{1}$ by 10.

$$10x = 52.222\dots \rightarrow \textcircled{2}$$

Subtracting $\textcircled{1}$ from $\textcircled{2}$, we get

$$10x = 52.222\dots$$

$$\leftarrow x = 5.222\dots$$

$$9x = 47.000$$

$$x = \frac{47}{9}$$

$\therefore x = 5.\bar{2} = \frac{47}{9}$, which is in $\frac{p}{q}$ form.

(iii) Let $x = 0.404040\dots \rightarrow \textcircled{1}$

As there is two repeating digit after the decimal point, so multiplying both sides of Eq $\textcircled{1}$ by 100

$$100x = 40.404040 \rightarrow \textcircled{2}$$

Subtracting $\textcircled{1}$ from $\textcircled{2}$, we get.

$$100x = 40.404040\dots$$

$$x = 0.404040\dots$$

\leftarrow

$$99x = 40.000$$

$$99x = 40$$

$$x = \frac{40}{99}$$

$\therefore x = 0.404040\dots = \frac{40}{99}$, which is in $\frac{p}{q}$ form.

(iv) Let $x = 0.4\bar{7} = 0.4777\dots$

$$x = 0.4777\dots \rightarrow \textcircled{1}$$

There is one non-repeating digit after the decimal point, multiplying both sides of $\textcircled{1}$ by 10.

$$10x = 4.777\dots \rightarrow \textcircled{2}$$

As there is one repeating digit after the decimal point, multiplying both sides of $\textcircled{2}$ by 10.

$$100x = 47.777\dots \rightarrow \textcircled{3}$$

Subtracting $\textcircled{2}$ from $\textcircled{3}$, we get

$$100x = 47.777\dots$$

$$- 10x = 4.777\dots$$

(-)

$$\hline 90x = 43.000$$

$$90x = 43$$

$$x = \frac{43}{90}$$

$\therefore x = 0.4777\dots = \frac{43}{90}$, which is in $\frac{p}{q}$ form.

(v) $0.\overline{134}$

Let $x = 0.1343434\dots \rightarrow \textcircled{1}$

There is one non-repeating digit after the decimal point, multiplying both sides of $\textcircled{1}$ by 10.

$10x = 1.343434\dots \rightarrow \textcircled{2}$

As there are two repeating digits after the decimal point, so multiplying both sides of $\textcircled{2}$ by 100.

$1000x = 134.343434\dots \rightarrow \textcircled{3}$

Subtracting $\textcircled{2}$ from $\textcircled{3}$, we get

$$\begin{array}{r} 1000x = 134.343434\dots \\ 10x = 1.343434\dots \\ \hline \end{array}$$

$990x = 133.00000$

—IGNITE YOUR LEARNING SPARK—

$990x = 133$

$x = \frac{133}{990}$

$\therefore x = \frac{133}{990}$, which is in $\frac{p}{q}$ form.

(vi) Let $x = 0.\overline{001}$

$x = 0.001001\dots \rightarrow \textcircled{1}$

As there are three repeating digits after the decimal point, so multiplying both sides $\textcircled{1}$ by 1000.

$1000x = 1.001001\dots \rightarrow \textcircled{2}$

Subtracting $\textcircled{1}$ from $\textcircled{2}$, we get,

$$1000x = 1.001001 \dots$$

$$x = 0.001001 \dots$$

$$\begin{array}{r} (-) \\ \hline 999x = 1 \end{array}$$

$$x = \frac{1}{999}$$

$\therefore x = 0.\overline{001} = \frac{1}{999}$, which is in $\frac{p}{q}$ form.

Solution (9)

(i) $\sqrt{23}$

Square root of 23 by long division method.

	4.79583
4	23.00000000
	16
87	700
	609
949	9100
	8541
9585	55900
	47925
95908	7,97500
	7,67,264
959163	30,23600
	28,77,489
	146111

$\therefore \sqrt{23} = 4.79583$, which has non-terminating and non-repeating decimal expansion.

So, it is an irrational number.

(ii) $\sqrt{225}$

Prime factorization of 225.

$$225 = 3 \times 3 \times 5 \times 5$$

$$225 = (3 \times 5)^2$$

$$\sqrt{225} = \sqrt{(3 \times 5)^2} = ((3 \times 5)^2)^{\frac{1}{2}}$$

$$\therefore \sqrt{225} = 3 \times 5 = 15.$$

$\sqrt{225} = 15$; which is a rational number.

3	225
3	75
5	25
5	5

(iii) 0.3796

Decimal Expansion of 0.3796 is terminating.

So, $0.3796 = \frac{3796}{10000}$ which is in $\frac{p}{q}$ form.

$\therefore 0.3796$ is a rational number.

(iv) $x = 7.478478 \dots \dots \rightarrow \textcircled{1}$

As there are three repeating digits after the decimal point, so multiplying both sides of $\textcircled{1}$ by 1000.

$$1000x = 7478.478478 \dots \dots \rightarrow \textcircled{2}$$

Subtracting $\textcircled{1}$ from $\textcircled{2}$, we get,

$$1000x = 7478.478478 \dots \dots$$

$$x = 7.478478 \dots \dots$$

(-)

$$999x = 7471.0$$

$$x = \frac{7471}{999}$$

$\therefore x = 7.478478\ldots = \frac{7471}{999}$, which
is in ' $\frac{p}{q}$ ' form.

So, $7.478478\ldots$ is a rational number.

(v) $1.101001000100001\ldots$

From the above decimal expansion, we observed that after decimal point, number of zeros between two consecutive ones are increasing. So, it is a non-terminating and non-repeating decimal expansion.

$\therefore 1.101001000100001\ldots$ is an irrational number.

(vi) $345.\overline{0456}$

Let $x = 345.0456456\ldots \rightarrow \textcircled{1}$

Multiplying by 10 on both sides of Eq. $\textcircled{1}$

$10x = 3450.456456\ldots \rightarrow \textcircled{2}$

As there are three repeating digits after the decimal point, so multiplying both sides of $\textcircled{2}$ by 1000.

$10000x = 3450456.456456\ldots \rightarrow \textcircled{3}$

$$\begin{array}{r} \textcircled{3} - \textcircled{2} \Rightarrow \\ 10000x = 3450456.456456\ldots \\ 10x = 345.456456\ldots \\ \hline (-) \end{array}$$

$$9990x = 3450111.0$$

$$\therefore 9990x = 3450111.$$

$$x = \frac{3450111}{9990},$$

which is in the form $\frac{p}{q}$

So, $345.04\overline{56}$ is a rational number.

Solution (10) :-

(i) Decimal Expansion of $\frac{1}{3}$ and $\frac{1}{2}$.

$$\therefore \frac{1}{3} = 0.333\dots$$

3	0.33
3	10
3	9
3	10
3	9

← remainder is repeating.

$$= 0.\overline{3}$$

$$\therefore \frac{1}{2} = 0.5$$

2	0.5
2	10
2	10
2	0

There are infinite rational numbers between $\frac{1}{3} (= 0.\overline{3})$ and $\frac{1}{2} (= 0.5)$.

One among them is $0.4040040004\dots$

(ii) $-\frac{2}{5}$ and $\frac{1}{2}$.

Decimal expansion of $-\frac{2}{5}$ and $\frac{1}{2}$.

$\therefore -\frac{2}{5} = -0.4$

$$\begin{array}{r|l} & 0.4 \\ 5 & 20 \\ & 20 \\ \hline & 0 \end{array}$$

$\therefore \frac{1}{2} = 0.5$

$$\begin{array}{r|l} & 0.5 \\ 2 & 10 \\ & 10 \\ \hline & 0 \end{array}$$

There are many irrational numbers between $-\frac{2}{5}$ and $\frac{1}{2}$. One among them is 0.1010010001...

(iii) 0 and 0.1

There are infinite irrational numbers between 0 and 0.1. One among them is

0.06006000600006.....

Solution (ii) :-

There are infinite irrational numbers between 2 and 3. Two among them are

2.0101001000100001.....

2.919119111911119.....

Solution (12) :-

Decimal Expansion of $\frac{4}{9}$ and $\frac{7}{11}$

$$\frac{4}{9} = 0.44 \dots \dots$$
$$= 0.\overline{4}$$

$$\begin{array}{r} 0.44 \\ 9 \overline{) 40} \\ \underline{36} \\ 40 \\ \underline{36} \\ 4 \end{array}$$

Remainder is repeating

$$\frac{7}{11} = 0.6363 \dots \dots$$
$$= 0.\overline{63}$$

$$\begin{array}{r} 0.63 \\ 11 \overline{) 70} \\ \underline{66} \\ 40 \\ \underline{33} \\ 7 \end{array}$$

Remainder is repeating.

there are infinite rational number between $\frac{4}{9} (= 0.\overline{4})$ and $\frac{7}{11} (= 0.\overline{63})$.

Two among them are $0.404004000400004 \dots \dots$,
 $0.515115111511115 \dots \dots$

Solution (13) :

$$\text{Value of } \sqrt{2} = 1.414 \dots \dots$$

$$\text{Value of } \sqrt{3} = 1.732 \dots \dots$$

there are many rational numbers between $\sqrt{2}$ and $\sqrt{3}$. One among them 1.6.

Finding Value of $\sqrt{2}$ and $\sqrt{3}$ by long division method.

$$\begin{array}{r}
 \sqrt{2} \\
 \hline
 1 \quad 414 \\
 \hline
 1 \quad 2.00 \ 00 \ 00 \\
 \quad 1 \\
 \hline
 24 \quad 100 \\
 \quad \quad 96 \\
 \hline
 281 \quad 400 \\
 \quad \quad 281 \\
 \hline
 2824 \quad 11900 \\
 \quad \quad \quad 11296 \\
 \hline
 \quad \quad \quad \quad 604
 \end{array}$$

stud flare
 $\sqrt{2} = 1.414$ LEARNING SPARK

$$\begin{array}{r}
 \sqrt{3} \\
 \hline
 1 \quad 732 \\
 \hline
 1 \quad 3.00 \ 00 \ 00 \\
 \quad 1 \\
 \hline
 27 \quad 200 \\
 \quad \quad 189 \\
 \hline
 343 \quad 1100 \\
 \quad \quad 1029 \\
 \hline
 3462 \quad 7100 \\
 \quad \quad \quad 6924 \\
 \hline
 \quad \quad \quad \quad 176
 \end{array}$$

$\therefore \sqrt{3} = 1.732 \dots\dots$

Solution (14):

$$2\sqrt{3} = \sqrt{2^2 \times 3} = \sqrt{4 \times 3} = \sqrt{4 \times 3}$$

$$\therefore 2\sqrt{3} = \sqrt{12}$$

We have, $12 < 12.25 < 12.96 < 15$

$$\Rightarrow \sqrt{12} < \sqrt{12.25} < \sqrt{12.96} < \sqrt{15}$$

$$\sqrt{12} < \sqrt{(3.5)^2} < \sqrt{(3.6)^2} < \sqrt{15}$$

$$\sqrt{12} < 3.5 < 3.6 < \sqrt{15}$$

\therefore 3.5 and 3.6 are two rational numbers between $\sqrt{12}$ and $\sqrt{15}$.

Solution (15):

We have, $5 < 6 < 7$.

$$\Rightarrow \sqrt{5} < \sqrt{6} < \sqrt{7}$$

$\therefore \sqrt{6}$ is an irrational number between $\sqrt{5}$ and $\sqrt{7}$.

Solution (16):

We have, $3 < 5 < 6 < 7$

$$\Rightarrow \sqrt{3} < \sqrt{5} < \sqrt{6} < \sqrt{7}$$

$\therefore \sqrt{5}$ and $\sqrt{6}$ are two irrational numbers between $\sqrt{3}$ and $\sqrt{7}$.

EXERCISE - 1.4

SOLUTION - 1

$$(i) \sqrt{45} - 3\sqrt{20} + 4\sqrt{5}$$

$$\begin{aligned} \text{Sol: } & \sqrt{9 \times 5} - 3\sqrt{4 \times 5} + 4\sqrt{5} \\ & = \sqrt{9}\sqrt{5} - 3\sqrt{4}\sqrt{5} + 4\sqrt{5} \\ & = 3\sqrt{5} - 3 \times 2\sqrt{5} + 4\sqrt{5} \\ & = 3\sqrt{5} - 6\sqrt{5} + 4\sqrt{5} \\ & = (3 - 6 + 4)\sqrt{5} \\ & = (7 - 6)\sqrt{5} \\ & = \sqrt{5} \end{aligned}$$

$$(ii) 3\sqrt{3} + 2\sqrt{27} + \frac{7}{\sqrt{3}}$$

$$\begin{aligned} \text{Sol: } & 3\sqrt{3} + 2\sqrt{9 \times 3} + \frac{7}{\sqrt{3}} \\ & = 3\sqrt{3} + 2 \times \sqrt{9}\sqrt{3} + \frac{7}{\sqrt{3}} \\ & = 3\sqrt{3} + 2 \times 3\sqrt{3} + \frac{7}{\sqrt{3}} \\ & = 3\sqrt{3} + 6\sqrt{3} + \frac{7}{\sqrt{3}} \\ & = (3 + 6)\sqrt{3} + \frac{7}{\sqrt{3}} \\ & = 9\sqrt{3} + \frac{7}{\sqrt{3}} \end{aligned}$$

Multiplying and Dividing by " $\sqrt{3}$ "

$$= 9\sqrt{3} + \frac{7}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}}$$

$$= 9\sqrt{3} + \frac{7\sqrt{3}}{3}$$

By cross multiplying

$$= \frac{3 \times 9\sqrt{3} + 7\sqrt{3}}{3}$$

$$= \frac{27\sqrt{3} + 7\sqrt{3}}{3}$$

$$= \frac{(27+7)\sqrt{3}}{3}$$

$$= \frac{34\sqrt{3}}{3}$$

stud flare
—IGNITE YOUR LEARNING SPARK—

(iii) $6\sqrt{5} \times 2\sqrt{5}$

Sol: $6 \times 2 \times \sqrt{5} \cdot \sqrt{5}$

$$= 12 \times (\sqrt{5})^2$$

$$= 12 \times 5$$

$$= 60$$

(iv) $8\sqrt{5} \div 2\sqrt{3}$

Sol: $\frac{8\sqrt{5}}{2\sqrt{3}} = \frac{4 \cdot 8 \cdot \sqrt{5} \sqrt{3}}{2\sqrt{3}}$

$$= 4\sqrt{5}$$

$$(V) \quad \frac{\sqrt{24}}{8} + \frac{\sqrt{54}}{9}$$

$$\text{Sol:} \quad \frac{\sqrt{6 \times 4}}{8} + \frac{\sqrt{9 \times 6}}{9}$$

$$= \frac{\sqrt{6} \cdot \sqrt{4}}{8} + \frac{\sqrt{9} \cdot \sqrt{6}}{9}$$

$$= \frac{2\sqrt{6}}{8} + \frac{3\sqrt{6}}{9}$$

$$= \frac{1 \cdot \sqrt{6}}{4} + \frac{1 \cdot \sqrt{6}}{3}$$

$$= \sqrt{6} \left[\frac{1}{4} + \frac{1}{3} \right]$$

$$= \sqrt{6} \left[\frac{3+4}{12} \right] \quad \because \text{LCM of 4 and 3}$$

$$= \frac{7\sqrt{6}}{12}$$

— NITE YOUR LEARNING SPARK —

$$(VI) \quad \frac{3}{\sqrt{8}} + \frac{1}{\sqrt{2}}$$

$$\text{Sol:} \quad \frac{3}{\sqrt{2 \times 4}} + \frac{1}{\sqrt{2}}$$

$$= \frac{3}{\sqrt{2} \cdot \sqrt{4}} + \frac{1}{\sqrt{2}}$$

$$= \frac{3}{2\sqrt{2}} + \frac{1}{\sqrt{2}}$$

$$= \frac{1}{\sqrt{2}} \left(\frac{3}{2} + 1 \right)$$

$$= \frac{1}{\sqrt{2}} \left(\frac{3+2}{2} \right)$$

\therefore LCM of 2 and 1
is 2

$$= \frac{1}{\sqrt{2}} \left(\frac{5}{2} \right)$$

$$= \frac{5}{2\sqrt{2}}$$

Multiply and Divide by " $\sqrt{2}$ "

$$= \frac{5}{2\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}}$$

$$= \frac{5\sqrt{2}}{2 \cdot \sqrt{2} \times \sqrt{2}}$$

$$= \frac{5\sqrt{2}}{2(\sqrt{2})^2} = \frac{5\sqrt{2}}{2 \cdot 2}$$

$$= \frac{5\sqrt{2}}{4}$$

stud flare

SOLUTION — IGNITE YOUR LEARNING SPARK —

(i) $(5 + \sqrt{7})(2 + \sqrt{5})$

Sol: $5 \times 2 + 5\sqrt{5} + 2\sqrt{7} + \sqrt{7} \cdot \sqrt{5}$

$$= 10 + 5\sqrt{5} + 2\sqrt{7} + \sqrt{7 \times 5}$$

$$= 10 + 5\sqrt{5} + 2\sqrt{7} + \sqrt{35}$$

(ii) $(5 + \sqrt{5})(5 - \sqrt{5})$

Sol: $(5)^2 - (\sqrt{5})^2$

$$= 25 - 5$$

$$= 20$$

$$(iii) (\sqrt{5} + \sqrt{2})^2$$

$$\text{Sol: } (\sqrt{5})^2 + (\sqrt{2})^2 + 2 \cdot \sqrt{5} \cdot \sqrt{2}$$

$$= 5 + 2 + 2\sqrt{5 \times 2}$$

$$= 5 + 2 + 2\sqrt{10}$$

$$= 7 + 2\sqrt{10}$$

$$(iv) (\sqrt{3} - \sqrt{7})^2$$

$$\text{Sol: } (\sqrt{3})^2 + (\sqrt{7})^2 - 2 \cdot \sqrt{3} \cdot \sqrt{7}$$

$$= 3 + 7 - 2\sqrt{3 \times 7}$$

$$= 10 - 2\sqrt{21}$$

$$(v) (\sqrt{2} + \sqrt{3})(\sqrt{5} + \sqrt{7})$$

$$\text{Sol: } \sqrt{2} \cdot \sqrt{5} + \sqrt{2} \cdot \sqrt{7} + \sqrt{3} \cdot \sqrt{5} + \sqrt{3} \cdot \sqrt{7}$$

$$= \sqrt{2 \times 5} + \sqrt{2 \times 7} + \sqrt{3 \times 5} + \sqrt{3 \times 7}$$

$$= \sqrt{10} + \sqrt{14} + \sqrt{15} + \sqrt{21}$$

$$(vi) (4 + \sqrt{5})(\sqrt{3} - \sqrt{7})$$

$$\text{Sol: } 4\sqrt{3} - 4\sqrt{7} + \sqrt{5} \cdot \sqrt{3} - \sqrt{5} \cdot \sqrt{7}$$

$$= 4\sqrt{3} - 4\sqrt{7} + \sqrt{5 \times 3} - \sqrt{5 \times 7}$$

$$= 4\sqrt{3} - 4\sqrt{7} + \sqrt{15} - \sqrt{35}$$

SOLUTION -3

~~~~~

(i)  $\sqrt{8} + \sqrt{50} + \sqrt{72} + \sqrt{98}$

Sol:  $\sqrt{4 \times 2} + \sqrt{25 \times 2} + \sqrt{36 \times 2} + \sqrt{49 \times 2}$

$$= \sqrt{4} \cdot \sqrt{2} + \sqrt{25} \cdot \sqrt{2} + \sqrt{36} \cdot \sqrt{2} + \sqrt{49} \cdot \sqrt{2}$$

$$= 2\sqrt{2} + 5\sqrt{2} + 6\sqrt{2} + 7\sqrt{2}$$

$$= (2+5+6+7)\sqrt{2}$$

$$= 20 \times \sqrt{2}$$

$$= 20 \times 1.414$$

$$= 28.28$$

(ii)  $3\sqrt{32} - 2\sqrt{50} + 4\sqrt{28} - 20\sqrt{18}$

—IGNITE YOUR LEARNING SPARK—

Sol:  $3\sqrt{16 \times 2} - 2\sqrt{25 \times 2} + 4\sqrt{64 \times 2} - 20\sqrt{9 \times 2}$

$$= 3 \times 4\sqrt{2} - 2 \times 5\sqrt{2} + 4 \times 8\sqrt{2} - 20 \times 3\sqrt{2}$$

$$= 12\sqrt{2} - 10\sqrt{2} + 32\sqrt{2} - 60\sqrt{2}$$

$$= (12 - 10 + 32 - 60)\sqrt{2}$$

$$= -26\sqrt{2}$$

$$= -26 \times 1.414$$

$$= -36.764$$

SOLUTION - 4

~~~~~

(i) $\sqrt{27} + \sqrt{75} + \sqrt{108} - \sqrt{243}$

Sol: $\sqrt{9 \times 3} + \sqrt{25 \times 3} + \sqrt{36 \times 3} - \sqrt{81 \times 3}$

$$= 3\sqrt{3} + 5\sqrt{3} + 6\sqrt{3} - 9\sqrt{3}$$

$$= (3+5+6-9)\sqrt{3}$$

$$= (14-9)\sqrt{3}$$

$$= 5 \times \sqrt{3}$$

$$= 5 \times 1.732$$

$$= 8.66$$

(ii) $3\sqrt{3/2} - 5\sqrt{12} - 3\sqrt{48} + 6\sqrt{75} + 7\sqrt{108}$

stud flare

IGNITE YOUR LEARNING SPARK

Sol: $5\sqrt{4 \times 3} - 3\sqrt{16 \times 3} + 6\sqrt{25 \times 3} + 7\sqrt{36 \times 3}$

$$= 5 \times 2\sqrt{3} - 3 \times 4\sqrt{3} + 6 \times 5\sqrt{3} + 7 \times 6\sqrt{3}$$

$$= 10\sqrt{3} - 12\sqrt{3} + 30\sqrt{3} + 42\sqrt{3}$$

$$= (10 - 12 + 30 + 42)\sqrt{3}$$

$$= (82 - 12)\sqrt{3}$$

$$= 70 \times \sqrt{3}$$

$$= 70 \times 1.732$$

$$= 121.24$$

SOLUTION - 5

(i) $\sqrt{\frac{4}{9}}$, $-\frac{3}{70}$, $\sqrt{\frac{7}{25}}$, $\sqrt{\frac{16}{5}}$

Sol: $\sqrt{\frac{4}{9}} = \frac{\sqrt{4}}{\sqrt{9}}$

$= \frac{2}{3}$. It is in the form of $\frac{p}{q}$

and p, q are integers

Therefore $\sqrt{\frac{4}{9}}$ is a rational number.

$\Rightarrow -\frac{3}{70}$ is a rational number

$\Rightarrow \sqrt{\frac{7}{25}} = \frac{\sqrt{7}}{\sqrt{25}}$

$= \frac{\sqrt{7}}{5}$. Since $\sqrt{7}$ is not an integer

Therefore, $\sqrt{\frac{7}{25}}$ is an irrational number.

$\Rightarrow \sqrt{\frac{16}{5}} = \frac{\sqrt{16}}{\sqrt{5}}$

$= \frac{4}{\sqrt{5}}$. Since $\sqrt{5}$ is not an integer

Therefore, $\sqrt{\frac{16}{5}}$ is an irrational number.

$$(ii) \quad -\sqrt{\frac{2}{49}}, \frac{3}{200}, \sqrt{\frac{25}{3}}, -\sqrt{\frac{49}{16}}$$

$$\text{Sol:} \quad -\sqrt{\frac{2}{49}} = -\frac{\sqrt{2}}{\sqrt{49}}$$

$$= -\frac{\sqrt{2}}{7} \quad \text{Since } \sqrt{2} \text{ is not an Integer}$$

Therefore, $-\sqrt{\frac{2}{49}}$ is an irrational number

$\Rightarrow \frac{3}{200}$ It is in the form of $\frac{p}{q}$
and p, q are Integers. So, it
is a rational number

$$\Rightarrow \sqrt{\frac{25}{3}} =$$

stud flare

—IGNITE YOUR LEARNING SPARK—

$$= \frac{5}{\sqrt{3}}$$

since $\sqrt{3}$ is not an Integer

Therefore, $\sqrt{\frac{25}{3}}$ is an irrational number

$$\Rightarrow -\sqrt{\frac{49}{16}} = -\frac{\sqrt{49}}{\sqrt{16}}$$

$$= -\frac{7}{4}$$

Therefore, $-\sqrt{\frac{49}{16}}$ is a rational number

SOLUTION-6

(i) $-3\sqrt{2}$

Sol: Since $\sqrt{2}$ is an irrational number

Therefore, $-3\sqrt{2}$ will change into non-terminating
non-recurring decimal

(ii) $\sqrt{\frac{256}{81}}$

Sol:- $\frac{\sqrt{256}}{\sqrt{81}} = \frac{16}{9}$

$= 1.777777$

Therefore, $\sqrt{\frac{256}{81}}$ will not change into non-terminating
non-recurring decimal

—IGNITE YOUR LEARNING SPARK—

(iii) $\sqrt{27 \times 16}$

Sol:- $\sqrt{27} \times \sqrt{16} = \sqrt{9 \times 3} \times 4$

$= 4 \times 3\sqrt{3}$

$= 12\sqrt{3}$

$\therefore \sqrt{3}$ is an irrational number

Therefore, $\sqrt{27 \times 16}$ will change into non-terminating
non-recurring decimal

(iv) $\sqrt{\frac{5}{36}}$

Sol:- $\frac{\sqrt{5}}{\sqrt{36}} = \frac{\sqrt{5}}{6}$

$\therefore \sqrt{5}$ is an irrational number

Therefore, $\sqrt{\frac{5}{36}}$ will change into non-terminating non-recurring
decimal

SOLUTION-7

(i) $3 - \sqrt{\frac{7}{25}}$

Sol: $3 - \frac{\sqrt{7}}{\sqrt{25}}$

$$= 3 - \frac{\sqrt{7}}{5}$$

$$= \frac{15 - \sqrt{7}}{5} \quad \therefore \sqrt{7} \text{ is an irrational number}$$

Therefore, $3 - \sqrt{\frac{7}{25}}$ is also an irrational number

(ii) $-\frac{2}{3} + \sqrt[3]{2}$

Sol: Since $\sqrt[3]{2}$ is an irrational number.

Therefore, $-\frac{2}{3} + \sqrt[3]{2}$ is also an irrational number

(NOTE: Sum of rational and irrational number is irrational)

(iii) $\frac{3}{\sqrt{3}}$

Sol: $\frac{\sqrt{3} \times \sqrt{3}}{\sqrt{3}} = \sqrt{3}$

Since $\sqrt{3}$ is an irrational number

Therefore, $\frac{3}{\sqrt{3}}$ is also an irrational number.

(iv) $-\frac{2}{7} \sqrt[3]{5}$

Sol: Since $\sqrt[3]{5}$ is an irrational number

Therefore, $-\frac{2}{7} \sqrt[3]{5}$ is also an irrational number.

(NOTE: Product of rational and irrational number is irrational)

(v) $(2-\sqrt{3})(2+\sqrt{3})$

Sol: $2 \times 2 + 2\sqrt{3} - 2\sqrt{3} - (\sqrt{3})^2$

$= 4 - 3$

$= 1$

Therefore, $(2-\sqrt{3})(2+\sqrt{3})$ is a rational number

(vi) $(3+\sqrt{5})^2$

Sol: $(3)^2 + (\sqrt{5})^2 + 2 \times 3 \times \sqrt{5}$

$= 9 + 5 + 6\sqrt{5}$

$= 14 + 6\sqrt{5}$

Since $\sqrt{5}$ is an irrational number

$(3+\sqrt{5})^2$ is also an irrational number

$$(VII), \left(\frac{2}{5}\sqrt{7}\right)^2$$

$$\text{Sol: } \left(\frac{2}{5}\right)^2 \cdot (\sqrt{7})^2$$

$$= \frac{4}{25} \times 7$$

$$= \frac{28}{25}$$

Therefore, $\left(\frac{2}{5}\sqrt{7}\right)^2$ is a rational number

$$(VIII), (3-\sqrt{6})^2$$

$$\text{Sol: } (3)^2 + (\sqrt{6})^2 - 2 \times 3 \times \sqrt{6}$$

$$= 9 + 6 - 6\sqrt{6}$$

$$= 15 - 6\sqrt{6}$$

Since $\sqrt{6}$ is an irrational number

$(3-\sqrt{6})^2$ is also an irrational number

SOLUTION - 8 :

$$(i) \sqrt[3]{2}$$

Sol: Suppose that $\sqrt[3]{2} = \frac{p}{q}$, where p, q are

integers, $q > 0$, p and q have no common factors (except 1)

$$2 = \left[\frac{p}{q}\right]^3$$

$$p^3 = 2q^3 \quad \rightarrow \textcircled{1}$$

As 2 divides $2q^3 \Rightarrow 2$ divides p^3

$\Rightarrow 2$ divides p

Let $p = 2k$, where k is an integer

Substituting this value of 'p' in (1), we get

$$(2k)^3 = 2q^3$$

$$8k^3 = 2q^3$$

$$4k^3 = q^3$$

As 2 divides $4k^3 \Rightarrow 2$ divides q^3

$\Rightarrow 2$ divides q

Thus p and q have a common factor "2".

This contradicts that p and q have no common factor (except 1)

Therefore, $\sqrt[3]{2}$ is an irrational number

(ii) $\sqrt[3]{3}$

Sol: Suppose that $\sqrt[3]{3} = \frac{p}{q}$, where p, q are integers

$q > 0$, p and q have no common factors

(except 1)

$$3 = \left(\frac{p}{q}\right)^3$$

$$p^3 = 3q^3 \rightarrow (1)$$

As 3 divides $3q^3 \Rightarrow 3$ divides p^3

$\Rightarrow 3$ divides p

Let $P = 3K$, where K is an Integer

Substituting this value of 'P' in ①, we get

$$(3K)^3 = 3q^3$$

$$27K^3 = 3q^3$$

$$9K^3 = q^3$$

As 3 divides $9K^3 \Rightarrow 3$ divides q^3

$\Rightarrow 3$ divide q

Thus P and q have a common factor "3"

This contradicts that P and q have no common factor (except 1)

Therefore, $\sqrt[3]{3}$ is an Irrational number.

stud flare

—IGNITE YOUR LEARNING SPARK—

(iii) $\sqrt[4]{5}$

Sol: Suppose that $\sqrt[4]{5} = \frac{p}{q}$, where p, q are Integers
 $q > 0$, p and q have no common factors
(except 1)

$$5 = \left(\frac{p}{q}\right)^4$$

$$p^4 = 5q^4 \rightarrow \text{①}$$

As 5 divides $5q^4 \Rightarrow 5$ divides p^4

$\Rightarrow 5$ divides p

Let $p = 5K$, where K is an Integer

Substituting this value of 'p' in ①, we get

$$(5K)^4 = 5q^4$$

$$625K^4 = 5q^4$$

$$125K^4 = q^4$$

As 5 divides $125K^4 \Rightarrow 5$ divides q^4

$\Rightarrow 5$ divides q

Thus p and q have a common factor "5"

This contradicts that p and q have no common factors (except 1)

Therefore, $\sqrt[4]{5}$ is an irrational number.

SOLUTION-9

IGNITE YOUR LEARNING SPARK

(i) $2\sqrt{3}$, $\frac{3}{\sqrt{2}}$, $-\sqrt{7}$, $\sqrt{15}$

Sol: $\sqrt{4 \times 3} = \sqrt{12}$

$$\frac{3}{\sqrt{2}} = \sqrt{\frac{9}{2}}$$

$$= \sqrt{4.5}$$

$$\therefore \sqrt{12}, \sqrt{4.5}, -\sqrt{7}, \sqrt{15}$$

$$-\sqrt{7} < \sqrt{4.5} < \sqrt{12} < \sqrt{15}$$

The greatest real number is $\sqrt{15}$

The smallest real number is $-\sqrt{7}$

$$(ii) -3\sqrt{2}, \frac{9}{\sqrt{5}}, -4, \frac{4}{3}\sqrt{5}, \frac{3}{2}\sqrt{3}$$

$$\text{Sol: } -3\sqrt{2} = -\sqrt{9 \times 2}$$

$$= -\sqrt{18}$$

$$\frac{9}{\sqrt{5}} = \sqrt{\frac{81}{5}}$$

$$= \sqrt{16.2}$$

$$-4 = -\sqrt{16}$$

$$\frac{4}{3}\sqrt{5} = \sqrt{\frac{16 \times 5}{9}}$$

$$= \sqrt{\frac{80}{9}}$$

$$= \sqrt{8.89}$$

$$\frac{3}{2}\sqrt{3} = \sqrt{\frac{9 \times 3}{4}}$$

$$= \sqrt{\frac{27}{4}}$$

$$= \sqrt{6.75}$$

$$\therefore -\sqrt{18}, \sqrt{16.2}, \sqrt{8.89}, -\sqrt{16}, \sqrt{6.75}$$

$$-3\sqrt{2} < -4 < \frac{3}{2}\sqrt{3} < \frac{4}{3}\sqrt{5} < \frac{9}{\sqrt{5}}$$

The greatest real number is $\frac{9}{\sqrt{5}}$

The smallest real number is $-3\sqrt{2}$

SOLUTION - 10

~~~~~

(i)  $3\sqrt{2}$ ,  $2\sqrt{3}$ ,  $\sqrt{15}$ , 4

Sol: Write all the numbers as square roots under one radical

$$3\sqrt{2} = \sqrt{9} \times \sqrt{2} = \sqrt{18}$$

$$2\sqrt{3} = \sqrt{4} \times \sqrt{3} = \sqrt{12}$$

$$\sqrt{15} = \sqrt{15}$$

$$4 = \sqrt{16}$$

Since  $12 < 15 < 16 < 18$

$$\Rightarrow \sqrt{12} < \sqrt{15} < \sqrt{16} < \sqrt{18}$$

$$\Rightarrow 2\sqrt{3} < \sqrt{15} < 4 < 3\sqrt{2}$$

Hence, the given numbers in ascending order are

$$2\sqrt{3}, \sqrt{15}, 4, 3\sqrt{2}$$

(ii)  $3\sqrt{2}$ ,  $2\sqrt{8}$ , 4,  $\sqrt{50}$ ,  $4\sqrt{3}$

Sol: Write all the numbers as square roots under one radical

$$3\sqrt{2} = \sqrt{9} \times \sqrt{2} = \sqrt{18}$$

$$2\sqrt{8} = \sqrt{4} \times \sqrt{8} = \sqrt{32}$$

$$4 = \sqrt{16}$$

$$\sqrt{50} = \sqrt{50}$$

$$4\sqrt{3} = \sqrt{16} \times \sqrt{3} = \sqrt{48}$$

Since  $16 < 18 < 32 < 48 < 50$

$$\Rightarrow \sqrt{16} < \sqrt{18} < \sqrt{32} < \sqrt{48} < \sqrt{50}$$

$$\Rightarrow 4 < 3\sqrt{2} < 2\sqrt{8} < 4\sqrt{3} < \sqrt{50}$$

Hence, the given numbers in ascending orders are

$$4, 3\sqrt{2}, 2\sqrt{8}, 4\sqrt{3}, \sqrt{50}$$

SOLUTION - 11

(i)  $\frac{9}{\sqrt{2}}, \frac{3}{2}\sqrt{5}, 4\sqrt{3}, 3\sqrt{\frac{6}{5}}$

Sol: Write all the numbers as square roots  
under one radical

$$\frac{9}{\sqrt{2}} = \sqrt{\frac{81}{2}} = \sqrt{40.5}$$

$$\frac{3}{2}\sqrt{5} = \sqrt{\frac{9}{4} \times 5} = \sqrt{\frac{45}{4}} = \sqrt{11.25}$$

$$4\sqrt{3} = \sqrt{16 \times 3} = \sqrt{48}$$

$$3\sqrt{\frac{6}{5}} = \sqrt{9 \times \frac{6}{5}} = \sqrt{\frac{54}{5}} = \sqrt{10.8}$$

Since  $48 > 40.5 > 11.25 > 10.8$

$$\Rightarrow \sqrt{48} > \sqrt{40.5} > \sqrt{11.25} > \sqrt{10.8}$$

$$\Rightarrow 4\sqrt{3} > \frac{9}{\sqrt{2}} > \frac{3}{2}\sqrt{5} > 3\sqrt{\frac{6}{5}}$$

Hence, the given numbers in descending orders

$$\text{are } 4\sqrt{3}, \frac{9}{\sqrt{2}}, \frac{3}{2}\sqrt{5}, 3\sqrt{\frac{6}{5}}$$

(ii)  $\frac{5}{\sqrt{3}}$ ,  $\frac{7}{3}\sqrt{2}$ ,  $-\sqrt{3}$ ,  $3\sqrt{5}$ ,  $2\sqrt{7}$

Sol: Write all the numbers as square roots under one radical

$$\frac{5}{\sqrt{3}} = \sqrt{\frac{25}{3}} = \sqrt{8.33}$$

$$\frac{7}{3}\sqrt{2} = \sqrt{\frac{49}{9}} \times \sqrt{2} = \sqrt{\frac{98}{9}} = \sqrt{10.89}$$

$$-\sqrt{3} = -\sqrt{3}$$

$$3\sqrt{5} = \sqrt{9} \times \sqrt{5} = \sqrt{45}$$

$$2\sqrt{7} = \sqrt{4} \times \sqrt{7} = \sqrt{28}$$

Since  $45 > 28 > 10.89 > 8.33 > -3$

$$\Rightarrow \sqrt{45} > \sqrt{28} > \sqrt{10.89} > \sqrt{8.33} > -\sqrt{3}$$

$$\Rightarrow 3\sqrt{5} > 2\sqrt{7} > \frac{7}{3}\sqrt{2} > \frac{5}{\sqrt{3}} > -\sqrt{3}$$

Hence, the given numbers in descending order

are  $3\sqrt{5}$ ,  $2\sqrt{7}$ ,  $\frac{7}{3}\sqrt{2}$ ,  $\frac{5}{\sqrt{3}}$ ,  $-\sqrt{3}$

SOLUTION - 12

(1)  $\sqrt[3]{2}$ ,  $\sqrt{3}$ ,  $\sqrt[6]{5}$

Sol: L.C.M. of 2, 3 and 6 is 6

$$\sqrt[3]{2} = 2^{\frac{1}{3}} = (2^2)^{\frac{1}{6}} = (4)^{\frac{1}{6}}$$

$$\sqrt{3} = 3^{\frac{1}{2}} = (3^3)^{\frac{1}{6}} = (27)^{\frac{1}{6}}$$

$$\sqrt[6]{5} = 5^{\frac{1}{6}} = (5^1)^{\frac{1}{6}} = (5)^{\frac{1}{6}}$$

Since  $4 < 5 < 27$

$$\Rightarrow (4)^{\frac{1}{6}} < (5)^{\frac{1}{6}} < (27)^{\frac{1}{6}}$$

$$\Rightarrow \sqrt[3]{2} < \sqrt[6]{5} < \sqrt{3}$$

Hence, the given numbers in ascending

order are  $\sqrt[3]{2}$ ,  $\sqrt[6]{5}$ ,  $\sqrt{3}$

## EXERCISE - 1.5

### SOLUTION - 1

(i)  $\frac{3}{4\sqrt{5}}$

Sol:  $\frac{3}{4\sqrt{5}} \times \frac{4\sqrt{5}}{4\sqrt{5}} = \frac{12\sqrt{5}}{16 \cdot (\sqrt{5})^2}$

$$= \frac{12\sqrt{5}}{16 \times 5}$$
$$= \frac{3\sqrt{5}}{20}$$

(ii)  $\frac{5\sqrt{7}}{\sqrt{3}}$

Sol:  $\frac{5\sqrt{7}}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{5\sqrt{7 \times 3}}{(\sqrt{3})^2}$

—IGNITE YOUR LEARNING SPARK—

$$= \frac{5\sqrt{21}}{3}$$

(iii)  $\frac{3}{4-\sqrt{7}}$

Sol:  $\frac{3}{4-\sqrt{7}} \times \frac{4+\sqrt{7}}{4+\sqrt{7}} = \frac{3(4+\sqrt{7})}{(4-\sqrt{7})(4+\sqrt{7})}$

$$= \frac{12 + 3\sqrt{7}}{16 - (\sqrt{7})^2}$$
$$= \frac{12 + 3\sqrt{7}}{16 - 7}$$
$$= \frac{12 + 3\sqrt{7}}{9}$$

$$= \frac{3(4 + 3\sqrt{7})}{9}$$

$$= \frac{4 + \sqrt{7}}{3}$$

(iv)  $\frac{17}{3\sqrt{2} + 1}$

Sol:  $\frac{17}{3\sqrt{2} + 1} \times \frac{3\sqrt{2} - 1}{3\sqrt{2} - 1} = \frac{17(3\sqrt{2} - 1)}{(3\sqrt{2} + 1)(3\sqrt{2} - 1)}$

$$= \frac{51\sqrt{2} - 17}{(3\sqrt{2})^2 - (1)^2}$$

$$= \frac{17(3\sqrt{2} - 1)}{9 \times 2 - 1}$$

$$= \frac{17(3\sqrt{2} - 1)}{18 - 1}$$

$$= \frac{17(3\sqrt{2} - 1)}{17}$$

$$= 3\sqrt{2} - 1$$

(v)  $\frac{16}{\sqrt{41} - 5}$

Sol:  $\frac{16}{\sqrt{41} - 5} \times \frac{\sqrt{41} + 5}{\sqrt{41} + 5} = \frac{16(\sqrt{41} + 5)}{(\sqrt{41})^2 - (5)^2}$

$$= \frac{16(\sqrt{41} + 5)}{41 - 25}$$

$$= \frac{16(\sqrt{41} + 5)}{16}$$

$$= \sqrt{41} + 5$$



$$(vi) \frac{1}{\sqrt{7}-\sqrt{6}}$$

$$\begin{aligned} \text{Sol: } \frac{1}{\sqrt{7}-\sqrt{6}} \times \frac{\sqrt{7}+\sqrt{6}}{\sqrt{7}+\sqrt{6}} &= \frac{\sqrt{7}+\sqrt{6}}{(\sqrt{7})^2 - (\sqrt{6})^2} \\ &= \frac{\sqrt{7}+\sqrt{6}}{7-6} \\ &= \sqrt{7}+\sqrt{6} \end{aligned}$$

$$(vii) \frac{1}{\sqrt{5}+\sqrt{2}}$$

$$\begin{aligned} \text{Sol: } \frac{1}{\sqrt{5}+\sqrt{2}} \times \frac{\sqrt{5}-\sqrt{2}}{\sqrt{5}-\sqrt{2}} &= \frac{\sqrt{5}-\sqrt{2}}{(\sqrt{5})^2 - (\sqrt{2})^2} \end{aligned}$$

stud flare

—IGNITE YOUR LEARNING SPARK—

$$(viii) \frac{\sqrt{2}+\sqrt{3}}{\sqrt{2}-\sqrt{3}}$$

$$\begin{aligned} \text{Sol: } \frac{\sqrt{2}+\sqrt{3}}{\sqrt{2}-\sqrt{3}} \times \frac{\sqrt{2}+\sqrt{3}}{\sqrt{2}+\sqrt{3}} &= \frac{(\sqrt{2}+\sqrt{3})^2}{(\sqrt{2})^2 - (\sqrt{3})^2} \\ &= \frac{(\sqrt{2})^2 + (\sqrt{3})^2 + 2 \cdot \sqrt{2} \cdot \sqrt{3}}{2-3} \\ &= \frac{2+3+2\sqrt{6}}{-1} \\ &= -(5+2\sqrt{6}) \\ &= -5-2\sqrt{6} \end{aligned}$$

SOLUTION-2

(i)  $\frac{7+3\sqrt{5}}{7-3\sqrt{5}}$

Sol:-  $\frac{7+3\sqrt{5}}{7-3\sqrt{5}} \times \frac{7+3\sqrt{5}}{7+3\sqrt{5}} = \frac{(7+3\sqrt{5})^2}{(7)^2 - (3\sqrt{5})^2}$

$$= \frac{(7)^2 + (3\sqrt{5})^2 + 2 \times 7 \times 3\sqrt{5}}{49 - 9 \times 5}$$
$$= \frac{49 + 45 + 14 \times 3\sqrt{5}}{49 - 45}$$
$$= \frac{94 + 42\sqrt{5}}{4}$$
$$= \frac{2(47 + 21\sqrt{5})}{4}$$
$$= \frac{47 + 21\sqrt{5}}{2}$$

stud flare  
—IGNITE YOUR LEARNING SPARK—

(ii)  $\frac{3-2\sqrt{2}}{3+2\sqrt{2}}$

Sol:-  $\frac{3-2\sqrt{2}}{3+2\sqrt{2}} \times \frac{3-2\sqrt{2}}{3-2\sqrt{2}} = \frac{(3-2\sqrt{2})^2}{(3)^2 - (2\sqrt{2})^2}$

$$= \frac{(3)^2 + (2\sqrt{2})^2 - 2 \times 3 \times 2\sqrt{2}}{9 - 4 \times 2}$$
$$= \frac{9 + 8 - 12\sqrt{2}}{9 - 8}$$
$$= \frac{17 - 12\sqrt{2}}{1}$$
$$= 17 - 12\sqrt{2}$$

$$(ii) \frac{5-3\sqrt{14}}{7+2\sqrt{14}}$$

$$\text{Sol: } \frac{5-3\sqrt{14}}{7+2\sqrt{14}} \times \frac{7-2\sqrt{14}}{7-2\sqrt{14}}$$

$$= \frac{5 \times 7 - 5 \times 2\sqrt{14} - 7 \times 3\sqrt{14} + 2 \times 3 \times \sqrt{14} \times \sqrt{14}}{(7)^2 - (2\sqrt{14})^2}$$

$$= \frac{35 - 10\sqrt{14} - 21\sqrt{14} + 6 \times 14}{49 - 4 \times 14}$$

$$= \frac{35 - 31\sqrt{14} + 84}{49 - 56}$$

$$= \frac{119 - 31\sqrt{14}}{-7}$$

$$= \frac{31\sqrt{14} - 119}{7}$$

SOLUTION - 3

$$(i) \frac{7\sqrt{3}}{\sqrt{10}+\sqrt{3}} - \frac{2\sqrt{5}}{\sqrt{6}+\sqrt{5}} - \frac{3\sqrt{2}}{\sqrt{15}+3\sqrt{2}}$$

$$\text{Sol: } \frac{7\sqrt{3}}{\sqrt{10}+\sqrt{3}} = \frac{7\sqrt{3}}{\sqrt{10}+\sqrt{3}} \times \frac{\sqrt{10}-\sqrt{3}}{\sqrt{10}-\sqrt{3}}$$

$$= \frac{7\sqrt{30} - 7 \times 3}{(\sqrt{10})^2 - (\sqrt{3})^2}$$

$$= \frac{7\sqrt{30} - 21}{10 - 3}$$

$$= \frac{7(\sqrt{30} - 3)}{7} = \sqrt{30} - 3$$

$$\begin{aligned} \frac{2\sqrt{5}}{\sqrt{6}+\sqrt{5}} &= \frac{2\sqrt{5}}{\sqrt{6}+\sqrt{5}} \times \frac{\sqrt{6}-\sqrt{5}}{\sqrt{6}-\sqrt{5}} \\ &= \frac{2\sqrt{30} - 2 \times 5}{(\sqrt{6})^2 - (\sqrt{5})^2} \\ &= \frac{2\sqrt{30} - 10}{6-5} \\ &= 2\sqrt{30} - 10 \end{aligned}$$

$$\begin{aligned} \frac{3\sqrt{2}}{\sqrt{15}+3\sqrt{2}} &= \frac{3\sqrt{2}}{\sqrt{15}+3\sqrt{2}} \times \frac{\sqrt{15}-3\sqrt{2}}{\sqrt{15}-3\sqrt{2}} \\ &= \frac{3\sqrt{30} - 9 \times 2}{(\sqrt{15})^2 - (3\sqrt{2})^2} \\ &= \frac{3\sqrt{30} - 18}{15 - 18} \\ &= \frac{-3(\sqrt{30} - 6)}{-2} \\ &= -\sqrt{30} + 6 \end{aligned}$$

$$\begin{aligned} \therefore \frac{7\sqrt{3}}{\sqrt{10}+\sqrt{3}} - \frac{2\sqrt{5}}{\sqrt{6}+\sqrt{5}} - \frac{3\sqrt{2}}{\sqrt{15}+3\sqrt{2}} \\ &= \sqrt{30} - 3 - (2\sqrt{30} - 10) - (-\sqrt{30} + 6) \\ &= \sqrt{30} - 3 - 2\sqrt{30} + 10 + \sqrt{30} - 6 \\ &= \cancel{\sqrt{30}} - \cancel{3} - \cancel{2\sqrt{30}} + 10 - 9 + 2\sqrt{30} - 2\sqrt{30} \\ &= \cancel{\sqrt{30}} - \cancel{2\sqrt{30}} \quad \downarrow \\ &= 1 \end{aligned}$$

### SOLUTION -4

$$(i) \frac{3-\sqrt{5}}{3+2\sqrt{5}} = -\frac{19}{11} + a\sqrt{5}$$

$$\begin{aligned}\text{Sol: } \frac{3-\sqrt{5}}{3+2\sqrt{5}} \times \frac{3-2\sqrt{5}}{3-2\sqrt{5}} &= \frac{(3-\sqrt{5})(3-2\sqrt{5})}{(3)^2 - (2\sqrt{5})^2} \\ &= \frac{9 - 6\sqrt{5} - 3\sqrt{5} + 2(5)}{9 - 4(5)} \\ &= \frac{19 - 9\sqrt{5}}{9 - 20} \\ &= \frac{+19 - 9\sqrt{5}}{-11}\end{aligned}$$

stud flare

—IGNITE YOUR LEARNING SPARK—

$$\therefore -\frac{19}{11} + a\sqrt{5} = -\frac{19}{11} + \frac{9}{11}\sqrt{5}$$

$$\Rightarrow a = \frac{9}{11}$$

$$(ii) \frac{\sqrt{2} + \sqrt{3}}{3\sqrt{2} - 2\sqrt{3}} = a - b\sqrt{6}$$

$$\begin{aligned}\text{Sol: } \frac{\sqrt{2} + \sqrt{3}}{3\sqrt{2} - 2\sqrt{3}} \times \frac{3\sqrt{2} + 2\sqrt{3}}{3\sqrt{2} + 2\sqrt{3}} &= \frac{(\sqrt{2} + \sqrt{3})(3\sqrt{2} + 2\sqrt{3})}{(3\sqrt{2})^2 - (2\sqrt{3})^2} \\ &= \frac{3(2) + 2\sqrt{6} + 3\sqrt{6} + 2(3)}{9(2) - 4(3)} \\ &= \frac{6 + 5\sqrt{6} + 6}{18 - 12}\end{aligned}$$

$$= \frac{12 + 5\sqrt{6}}{6}$$

$$= 2 + \frac{5}{6}\sqrt{6}$$

$$= 2 - \left(-\frac{5}{6}\right)\sqrt{6}$$

$$\therefore a - b\sqrt{6} = 2 - \left(-\frac{5}{6}\right)\sqrt{6}$$

$$\Rightarrow a = 2 \quad ; \quad b = -\frac{5}{6}$$

$$(iii) \quad \frac{7+\sqrt{5}}{7-\sqrt{5}} - \frac{7-\sqrt{5}}{7+\sqrt{5}} = a + \frac{7}{11}b\sqrt{5}$$

$$\text{Sol: } \frac{7+\sqrt{5}}{7-\sqrt{5}} \times \frac{7+\sqrt{5}}{7+\sqrt{5}} = \frac{(7+\sqrt{5})^2}{(7)^2 - (\sqrt{5})^2}$$

$$= \frac{49 + (\sqrt{5})^2 + 2 \cdot 7 \cdot \sqrt{5}}{49 - 5}$$

$$= \frac{49 + 5 + 14\sqrt{5}}{44}$$

$$\frac{7-\sqrt{5}}{7+\sqrt{5}} \times \frac{7-\sqrt{5}}{7-\sqrt{5}} = \frac{(7-\sqrt{5})^2}{(7)^2 - (\sqrt{5})^2}$$

$$= \frac{49 + (\sqrt{5})^2 - 2 \times 7 \times \sqrt{5}}{49 - 5}$$

$$= \frac{49 + 5 - 14\sqrt{5}}{44}$$

$$\therefore \frac{7+\sqrt{5}}{7-\sqrt{5}} - \frac{7-\sqrt{5}}{7+\sqrt{5}} = \frac{54 + 14\sqrt{5}}{44} - \frac{54 - 14\sqrt{5}}{44}$$

$$= \frac{54 + 14\sqrt{5} - 54 + 14\sqrt{5}}{44}$$

$$= \frac{7 \cdot 28\sqrt{5}}{11 \cdot 44}$$

$$= \frac{7}{11} \times \sqrt{5}$$

$$\therefore a + \frac{7}{11} b\sqrt{5} = \frac{7}{11} \sqrt{5}$$

$$\Rightarrow a=0 ; b=1$$

SOLUTION - 5 :

$$(i) \frac{7+3\sqrt{5}}{3+\sqrt{5}} - \frac{7-3\sqrt{5}}{3-\sqrt{5}} = p+q\sqrt{5}$$

Sol:  $\frac{7+3\sqrt{5}}{3+\sqrt{5}} \times \frac{3-\sqrt{5}}{3-\sqrt{5}} = \frac{(7+3\sqrt{5})(3-\sqrt{5})}{(3)^2 - (\sqrt{5})^2}$

$$= \frac{21 - 7\sqrt{5} + 9\sqrt{5} - 3(5)}{4}$$

stud flare  
— IGNITE YOUR LEARNING SPARK —

$$= \frac{21 + 2\sqrt{5} - 15}{4}$$

$$= \frac{6+2\sqrt{5}}{4}$$

$$\frac{7-3\sqrt{5}}{3-\sqrt{5}} \times \frac{3+\sqrt{5}}{3+\sqrt{5}} = \frac{(7-3\sqrt{5})(3+\sqrt{5})}{(3)^2 - (\sqrt{5})^2}$$

$$= \frac{21 + 7\sqrt{5} - 9\sqrt{5} - 3(5)}{9-5}$$

$$= \frac{21 - 2\sqrt{5} - 15}{4}$$

$$= \frac{6-2\sqrt{5}}{4}$$

$$\begin{aligned}
 \therefore \frac{7+3\sqrt{5}}{3+\sqrt{5}} - \frac{7-3\sqrt{5}}{3-\sqrt{5}} &= \frac{6+2\sqrt{5}}{4} - \frac{6-2\sqrt{5}}{4} \\
 &= \frac{6+2\sqrt{5} - 6 + 2\sqrt{5}}{4} \\
 &= \frac{4\sqrt{5}}{4} \\
 &= \sqrt{5}
 \end{aligned}$$

$$\therefore p+q\sqrt{5} = \sqrt{5}$$

$$\Rightarrow p=0 ; q=1$$

SOLUTION -6 :

(i)  $\frac{\sqrt{2}}{2+\sqrt{2}}$

Sol:  $\frac{\sqrt{2}}{2+\sqrt{2}} \times \frac{2-\sqrt{2}}{2-\sqrt{2}} = \frac{\sqrt{2}(2-\sqrt{2})}{(2)^2 - (\sqrt{2})^2}$

$$= \frac{2\sqrt{2} - 2}{4 - 2}$$

$$= \frac{2(\sqrt{2} - 1)}{2}$$

$$= \sqrt{2} - 1$$

$$= 1.414 - 1$$

$$= 0.414$$



$$(ii) \frac{1}{\sqrt{3}+\sqrt{2}}$$

$$\begin{aligned}\text{Sol: } \frac{1}{\sqrt{3}+\sqrt{2}} \times \frac{\sqrt{3}-\sqrt{2}}{\sqrt{3}-\sqrt{2}} &= \frac{\sqrt{3}-\sqrt{2}}{(\sqrt{3})^2 - (\sqrt{2})^2} \\ &= \frac{\sqrt{3}-\sqrt{2}}{3-2} \\ &= \sqrt{3}-\sqrt{2} \\ &= 1.732 - 1.414 \\ &= 0.318\end{aligned}$$

SOLUTION - 7 :

$$(i) a = 2 + \sqrt{3}$$

$$\text{Sol: } \frac{1}{a} = \frac{1}{2+\sqrt{3}} = \frac{1}{2+\sqrt{3}} \times \frac{2-\sqrt{3}}{2-\sqrt{3}}$$

$$\begin{aligned}&= \frac{2-\sqrt{3}}{(2)^2 - (\sqrt{3})^2} \\ &= \frac{2-\sqrt{3}}{4-3} \\ &= 2-\sqrt{3}\end{aligned}$$

$$\begin{aligned}\therefore a - \frac{1}{a} &= 2 + \sqrt{3} - (2 - \sqrt{3}) \\ &= 2 + \sqrt{3} - 2 + \sqrt{3} \\ &= 2\sqrt{3}\end{aligned}$$

SOLUTION - 8

(i)  $x = 1 - \sqrt{2}$

Sol: Given  $x = 1 - \sqrt{2}$

$$\begin{aligned}\therefore \frac{1}{x} &= \frac{1}{1 - \sqrt{2}} = \frac{1}{1 - \sqrt{2}} \times \frac{1 + \sqrt{2}}{1 + \sqrt{2}} \\ &= \frac{1 + \sqrt{2}}{(1)^2 - (\sqrt{2})^2} \\ &= \frac{1 + \sqrt{2}}{1 - 2} \\ &= -(1 + \sqrt{2})\end{aligned}$$

$$\begin{aligned}\therefore \left(x - \frac{1}{x}\right)^4 &= (1 - \sqrt{2} - (-1 - \sqrt{2}))^4 \\ &= (1 - \sqrt{2} + 1 + \sqrt{2})^4 \\ &= 16\end{aligned}$$

stud flare  
—IGNITE YOUR LEARNING SPARK—

SOLUTION - 9

(i)  $x = 5 - 2\sqrt{6}$

Sol: Given  $x = 5 - 2\sqrt{6}$

$$\begin{aligned}\therefore \frac{1}{x} &= \frac{1}{5 - 2\sqrt{6}} = \frac{1}{5 - 2\sqrt{6}} \times \frac{5 + 2\sqrt{6}}{5 + 2\sqrt{6}} \\ &= \frac{5 + 2\sqrt{6}}{(5)^2 - (2\sqrt{6})^2} \\ &= \frac{5 + 2\sqrt{6}}{25 - 24} \\ &= 5 + 2\sqrt{6}\end{aligned}$$

$$\begin{aligned}\therefore x + \frac{1}{x} &= (5 - 2\sqrt{6}) + (5 + 2\sqrt{6}) \\ &= 10\end{aligned}$$

We know that  $(x + \frac{1}{x})^2 = x^2 + \frac{1}{x^2} + 2$

$$\begin{aligned}\Rightarrow x^2 + \frac{1}{x^2} &= (x + \frac{1}{x})^2 - 2 \\ &= (10)^2 - 2 \\ &= 100 - 2 \\ &= 98\end{aligned}$$

SOLUTION - 10

(i)  $p = \frac{2 - \sqrt{5}}{2 + \sqrt{5}}$

$q = \frac{2 + \sqrt{5}}{2 - \sqrt{5}}$

stud flare

— IGNITE YOUR LEARNING SPARK —

$$\begin{aligned}\text{Sol: } p + q &= \frac{2 - \sqrt{5}}{2 + \sqrt{5}} + \frac{2 + \sqrt{5}}{2 - \sqrt{5}} \\ &= \frac{(2 - \sqrt{5})^2 + (2 + \sqrt{5})^2}{(2)^2 - (\sqrt{5})^2} \\ &= \frac{(4 + 5 - 4\sqrt{5}) + (4 + 5 + 4\sqrt{5})}{4 - 5} \\ &= \frac{18}{-1}\end{aligned}$$

$\therefore p + q = -18$

$$\begin{aligned}
 \text{(ii) } p - q &= \frac{2-\sqrt{5}}{2+\sqrt{5}} - \frac{2+\sqrt{5}}{2-\sqrt{5}} \\
 &= \frac{(2-\sqrt{5})^2 - (2+\sqrt{5})^2}{(2)^2 - (\sqrt{5})^2} \\
 &= \frac{(4+5-4\sqrt{5}) - (4+5+4\sqrt{5})}{4-5} \\
 &= \frac{9-4\sqrt{5} - 9-4\sqrt{5}}{-1} \\
 &= -\frac{8\sqrt{5}}{-1} \\
 &= 8\sqrt{5}
 \end{aligned}$$

(iii)  $p^2 + q^2$

stud flare

Sol: We know that IGNITE YOUR LEARNING SPARK

$$(p+q)^2 = p^2 + q^2 + 2pq$$

$$\therefore pq = \frac{2-\sqrt{5}}{2+\sqrt{5}} \times \frac{2+\sqrt{5}}{2-\sqrt{5}} = 1$$

$$\therefore p+q = -18$$

$$\begin{aligned}
 \Rightarrow p^2 + q^2 &= (p+q)^2 - 2pq \\
 &= (-18)^2 - 2 \times 1 \\
 &= 324 - 2 \\
 &= 322
 \end{aligned}$$

$$(iv) p^2 - q^2$$

$$\begin{aligned}\text{Sol:} \quad \therefore p^2 - q^2 &= (p+q)(p-q) \\ &= (-18)(8\sqrt{5}) \\ &= -144\sqrt{5}\end{aligned}$$

stud flare

—IGNITE YOUR LEARNING SPARK—